Colonoscopy Bowel Preparation Instructions

 \Box SPLIT Dose: Suprep, Moviprep, Clenpiq \Box SPLIT Dose: Golytely, Nulytely, Trilyte or Colyte

Online Instructions:

www.eehealth.org/services/gastrointestinal/patient-instructions

Endeavor Health Gastroenterology

Christopher A. Boutin, MD Michael Greenspan, MD Jason Kramer, MD Shubha Singh, MD George Stathopoulos, MD Atena Lodhi, MD Patrick J. Lynch, MD S. Dharan Kumar, MD Karen Ma, MD Allison Rzepczynski MD

•	Your appointment is on: (day of the week)	(date)	(arrival time)	a.m/p.m
•	You are scheduled to have your test done at:			
	□ Elmhurst Memorial Hospital: 155 E. Brush Hill Road	d. Park in Green Lot, East E	intrance. 2 nd floor	
	Endoscopy registration desk.			
	□ North Elmhurst Location: 755 N. York St. Park in ma	ain entrance and register at	front desk.	
	☐ Elmhurst Outpatient Surgery Center/Center for If you do not receive your arrival time the day before Health directly at 630.758.8888.			
	NOTE: It is the patient's responsibility to check with the insura as for any questions regarding benefits, & out-of-pocket costs. P			ormation.

READ ALL OF THESE INSTRUCTIONS SEVERAL DAYS BEFORE YOUR SCHEDULED TEST DATE

Diagnosis Code(s): __

STARTING FIVE DAYS BEFORE THE COLONOSCOPY:

- 1. Pick up your prescription prep from your pharmacy. Call your pharmacy directly for pick up time and cost.
- 2. Pick up a small pack of simethicone tablets (anti-gas chewable or soft gels such as Gas-X, Mylanta Gas, Maalox Anti-Gas, or similar products). This is over the counter/no prescription required.
- 3. Recommended diet: meats, fish, breads, pasta, cereal, rice, mashed potatoes, bananas, applesauce, eggs, and canned peaches/pears.
- 4. DO NOT EAT: Fruits and vegetables, corn, nuts, seeds and popcorn.

ONE DAY BEFORE THE COLONOSCOPY:

- 1. **You may have BREAKFAST BEFORE 10 AM** (apple sauce, eggs, OR a piece of white toast nothing with nuts or seeds).
- 2. FOR LUNCH & DINNER, NO SOLID FOODS ONLY DRINK CLEAR LIQUIDS (See chart on the back).
- 3. BOWEL PREP:

Procedure Code(s):

- START DRINKING PREP AT: 5:00 PM. Complete FIRST HALF by 8:00 PM.
- 4. At 9:00PM: Take 2 simethicone anti-gas chewable or soft gels with 8 ounces of clear liquid. The simethicone medication reduces bubble formation in your colon and improves exam quality.
 - i. If you feel nauseated or bloated when drinking, take a short break and walk around to help the liquid pass through your intestines.

DAY OF COLONOSCOPY:

- 1. NO SOLID FOODS!
- 2. BOWEL PREP:
 - a. Complete **SECOND HALF** of bowel prep.
 - b. **START** drinking prep **(6 Hours** prior to procedure time) AT: ___
 - c. **FINISH** drinking prep (3 Hours Prior to procedure time) BY:
 - d. Stool must be liquid and clear without solid material to proceed with a successful colonoscopy.
- 3. May have **CLEAR LIQUIDS** up to 3 hours before procedure time. If you fail to keep your stomach empty for three hours, your procedure may be **CANCELLED**.

TRANSPORTATION

Plan to have someone drive you home after the procedure. You <u>CANNOT</u> use public transportation (Uber, Lyft, Taxi). The procedure(s) involves sedation, and you will not be allowed to leave unaccompanied. <u>NO EXCEPTIONS.</u> Your procedure CANNOT be performed if you do not have a driver/adult to escort you home.

WHAT TO WEAR

Wear casual clothes. Please leave jewelry and valuables at home. You may wear your dentures and eyeglasses, however these will be

removed prior to the procedure. Please let the staff know that you are wearing dentures before the procedure.

WHAT TO BRING

A picture ID, proof of insurance, list of your medications and allergies, and your referral form (if your insurance requires it).

QUESTIONS/RESCHEDULING

If you have any questions, please do not hesitate to call us at 331-221-9003 (office). To cancel or reschedule, please contact the office.

AT LEAST 7 Business days prior to your scheduled procedure. Please see attached cancellation policy.



$\underline{MEDICATION(S)} \rightarrow \underline{OK} \ \underline{TO} \ \underline{TAKE} \ \underline{BEFORE} \ \underline{THE} \ \underline{PROCEDURE}$:

- Aspirin
- Statins & Seizure medications → OK to continue
- BETA BLOCKERS → OK to continue
- ACE INHIBITORS, ANGIOTENSIN II RECEPTOR BLOCKERS (ARBs) & COMBINATION DRUGS THAT CONTAIN AN $ARB \rightarrow OK$ to continue

MEDICATION(S) TO HOLD \rightarrow DO NOT TAKE BEFORE PROCEDURE (READ BELOW):

- Iron (ferrous sulfate/ ferrous gluconate), Herbal Supplements & Multivitamins HOLD 7 DAYS (1 WEEK)
 prior to the procedure.
- Ozempic, Wegovy, Mounjaro, Trulicity, Rybelsus, Victoza HOLD 7 DAYS (1 WEEK) prior to the procedure.
- Jardiance, Invokana, Farxiga, Steglatro, Brenzavvy HOLD 4 DAYS (96 hours) prior to the procedure.
- <u>ANOREXIANTS</u> (Weight Loss Medication i.e., <u>Phentermine/Vyvanse</u>) <u>HOLD 7 days</u> (1 WEEK) prior to the procedure.
- <u>DO NOT TAKE:</u> Any form of <u>Erectile Dysfunction</u> medications for 3 days (72 hours) prior to the procedure UNLESS prescribed for Pulmonary Hypertension.
- DO NOT TAKE: Any form of alcohol and recreational drugs for 1 DAY (24 hours) prior to the
 procedure.

PATIENTS ON BLOOD THINNERS (ANTICOAGULANTS)

- YOU are responsible for contacting your cardiologist or prescribing provider, regarding recommendations on holding these
 therapies prior to your scheduled procedure.
- o This should be done at least 2 weeks in advance, as some medications will need to be held 5 days prior to your procedure.
- *IF you forget to stop your blood thinner, your procedure WILL be cancelled and rescheduled*.

PATIENTS ON DIABETIC MEDICATIONS (ORAL/INJECTIONS)

- HOLD ORAL diabetic medication(s) the DAY BEFORE and DAY OF procedure OR per MD discretion. *This will be reviewed with you when scheduling procedure*!
- IF YOU TAKE INSULIN
 - <u>YOU</u> will need to contact your **endocrinologist** or **prescribing provider**, prior to your scheduled procedure for recommendations on adjustments to these therapies during the bowel prep process and the day of your procedure.
 - o This should be done at least TWO WEEKS in advance.
 - o Bring your insulin with you to the exam.
- *Adjustments to these therapies are to ensure your safety during the bowel prep process and procedure. Failure to contact your providers and make the necessary adjustments to your medications prior to your scheduled procedure date may result in the cancellation of your procedure*.
- Additional Instructions:

CLEAR LIQUID DIET

NOT ACCEPTABLE	ACCEPTABLE/ALLOWED	
 SOLID FOOD ANYTHING THAT IS RED, BLUE, PURPLE DAIRY/MILK PRODUCTS/HONEY ALCOHOL TOMATO JUICE OR ANY FRUIT JUICE WITH PULP 	 LEMON-LIME SODA (SPRITE, GINGER-ALE, 7UP, CLUB SODA) BROTH (VEGETABLE, CHICKEN, BEEF → LIQUID ONLY, NO SOLID FOOD!) WATER BLACK COFFEE/TEA (NO CREAMERS, DAIRY/MILK, HONEY) APPLE JUICE, WHITE GRAPE JUICE, LEMONADE POPSCILES, ITALIAN ICE (NOT RED, BLUE, PURPLE) JELL-O (NOT RED, BLUE, PURPLE OR MILK-BASED) GATORADE, PROPEL, POWER-ADE (NOT RED, BLUE, PURPLE) 	