🥭 Endeavor Health..

Colonoscopy Bowel Preparation Instructions

<u>SPLIT</u> Dose MiraLAX + Gatorade

Online Instructions: <u>www.eehealth.org/services/gastrointestinal/patie</u> <u>nt-instructions/</u>

Endeavor Health Gastroenterology

Christopher A. Boutin, MD Michael Greenspan, MD Jason Kramer, MD Shubha Singh, MD George Stathopoulos, MD Atena Lodhi, MD Patrick J. Lynch, MD S. Dharan Kumar, MD Karen Ma, MD Allison Rzepczynski, MD

• Your appointment is on: (day of the week) _____(date) _____(arrival time) _____a.m/p.m.

• You are scheduled to have your test done at:

🗆 Elmhurst Memorial Hospital: 155 E. Brush Hill Road. Park in Green Lot, East Entrance. 2nd floor Endoscopy registration desk.

□ North Elmhurst Location: 755 N. York St. Park in main entrance and register at front desk.

<u>Elmhurst Hospital/North Elmhurst Location:</u> If you do not receive your arrival time the day before your procedure by 5PM, please call 331-221-1167.

□ Elmhurst Outpatient Surgery Center/Center for Health: 1200 South York Rd. Park in Purple Lot. Go to 1st floor registration desk.

- If you do not receive your arrival time the day before your procedure by 3PM, please call the <u>Center for Health</u> directly at 630.758.88888.
- If scheduled at <u>Elmhurst Outpatient Surgery Center</u>: your transportation will need to remain at the facility for the <u>entire duration of</u> <u>the procedure</u>. If your ride is unable to stay on-site, there is a risk that your appointment may be canceled.

NOTE: It is the patient's responsibility to check with the insurance company to see if pre-certification is required, as well as for any questions regarding benefits, & out-of-pocket costs. See attached sheet for procedural and diagnostic code information.

Procedure Code(s): _

Diagnosis Code(s): _

READ ALL OF THESE INSTRUCTIONS SEVERAL DAYS BEFORE YOUR SCHEDULED TEST DATE

For this preparation, please purchase the following items <u>OVER THE COUNTER</u> items at your local pharmacy:

- 1. One (1) **238 g bottle** of MiraLAX
- 2. Two (2) Dulcolax **laxative** tablets.
- 3. 64 oz. Gatorade (<u>NOT</u> red, purple, blue)
- 4. Small pack of simethicone tablets (anti-gas chewable or soft gels such as Gas-X, Mylanta Gas, Maalox Anti-Gas, or similar products) **STARTING FIVE DAYS BEFORE THE COLONOSCOPY:**
 - 1. Pick up your bowel prep over the counter (as above) at your pharmacy, no prescription is required.
 - 2. Recommended diet: meats, fish, breads, pasta, cereal, rice, mashed potatoes, bananas, applesauce, eggs, and canned peaches/pears.

3. DO NOT EAT: fruits and vegetables, corn, nuts, seeds and popcorn.

ONE DAY BEFORE THE COLONOSCOPY:

- NO SOLID FOOD! Clear liquids only.
- Breakfast, Lunch, Dinner: Only drink clear liquids (See list on the back).
- BOWEL PREP:
 - Step 1: Beginning at 2:00 PM, take two dulcolax laxative tablets with 8 oz. of water. Swallow the tablets whole with a full glass of water.
 - Step 2: At 5:00 PM, mix the bottle of MiraLAX in <u>64 oz.</u> of Gatorade. Shake the solution until the MiraLAX is dissolved. Complete <u>FIRST HALF</u> by <u>8:00 PM</u>. Put the remaining solution in the refrigerator.
 - **Step 3:** At <u>**9:00PM**</u>: Take 2 simethicone anti-gas chewable or soft gels with 8 ounces of clear liquid. The simethicone medication reduces bubble formation in your colon and improves exam quality.
 - o If you feel nauseated or bloated when drinking, take a short break and walk around to help the liquid pass through your intestines.

DAY OF COLONOSCOPY:

NO SOLID FOODS!

BOWEL PREP:

- Complete **SECOND HALF** of **MiraLAX/Gatorade)** bowel prep.
- START DRINKING PREP (6 hours prior to procedure time) AT: ______
- FINISH DRINKING PREP (3 hours prior to procedure time) BY: ____
- Stool must be liquid and clear without solid material to proceed with a successful colonoscopy.
 - You may have <u>**CLEAR LIQUIDS</u>** up to **3 hours** before procedure time.</u>
- If you fail to keep your stomach empty for three hours, your procedure may be canceled.

TRANSPORTATION

Plan to have someone drive you home after the procedure. You <u>CANNOT</u> use public transportation (Uber, Lyft, Taxi). The procedure (s) involves sedation, and you will not be allowed to leave unaccompanied. <u>NO EXCEPTIONS.</u>

≡ Endeavor Health..

Wear casual clothes. Please leave jewelry and valuables at home. You may wear your dentures and eyeglasses, however these will be removed prior to the procedure. Please let the staff know that you are wearing dentures before the procedure.

WHAT TO BRING

A picture ID, proof of insurance, list of your medications and allergies, and your referral form (if your insurance requires it).

<u>QUESTIONS/RESCHEDULING</u>: If you have any questions, please do not hesitate to call us at **331-221-9003**. To cancel or reschedule please contact the office **AT LEAST 7 business days** prior to your scheduled procedure.

MEDICATIONS OK TO TAKE BEFORE THE PROCEDURE:

- You may take your medications as directed, unless advised otherwise (see below).
- Statins & Seizure medications
- Beta Blockers

MEDICATIONS TO STOP BEFORE PROCEDURE:

- ACE inhibitors, Angiotensin II receptor blockers (ARBs) & Combination Drugs that contain ARBs: Discontinue the <u>night</u> <u>before</u> OR the <u>day of</u> procedure.
- Iron (ferrous sulfate/ ferrous gluconate), Herbal Supplements & Multivitamins: Discontinue 2 weeks prior to procedure.
- Ozempic, Wegovy, Mounjaro, Trulicity, Rybelsus, Victoza, Saxenda: Discontinue 1 week prior to procedure.
- Jardiance, Invokana, Farxiga, Steglatro, Brenzavvy: Discontinue 4 days prior to the procedure.
- <u>ANOREXIANTS</u> (weight loss medication i.e., <u>Phentermine/Vyvanse)</u>: Discontinue 7 days prior to the procedure.
- <u>ADHD Medications</u>: Discontinue the day of procedure.
- <u>Diuretics</u>: Discontinue the day of procedure.
- <u>DO NOT TAKE</u>: Any form of <u>erectile dysfunction</u> medication for 3 days prior to the procedure *unless* taken for *Pulmonary Hypertension or Prostatic Hyperplasia*.
- DO NOT TAKE: Any form of alcohol and recreational drugs for 24 hours prior to the procedure.

PATIENTS ON BLOOD THINNERS (ANTICOAGULANTS)

- <u>YOU</u> are responsible for contacting your **cardiologist** or **prescribing provider**, regarding recommendations on holding these therapies prior to your scheduled procedure.
- This should be done **at least 2 weeks in advance**, as some medications will need to be held 5 days prior to your procedure.
- *IF you forget to stop your blood thinner, your procedure will be cancelled and rescheduled.

PATIENTS ON DIABETIC MEDICATIONS (ORAL/INJECTIONS)

- **HOLD** all oral diabetic medications the **<u>day before</u>** and **<u>day of</u> procedure**.
 - IF YOU TAKE INSULIN:
 - **YOU** will need to contact your **endocrinologist** &/or **prescribing provider**, prior to your scheduled procedure for recommendations on adjustments to these therapies during the bowel prep process and the day of your procedure.
 - This should be done at least two weeks in advance.
 - Bring your insulin with you to the exam.
- Additional Instructions:

CLEAR LIQUID DIET

NOT ACCEPTABLE	ACCEPTABLE/ALLOWED
 SOLID FOOD ANYTHING THAT IS RED, BLUE, PURPLE DAIRY/MILK PRODUCTS/HONEY ALCOHOL TOMATO JUICE OR ANY FRUIT JUICE WITH PULP 	 LEMON-LIME SODA (SPRITE, GINGER-ALE, 7UP, CLUB SODA) BROTH (VEGETABLE, CHICKEN, BEEF → LIQUID ONLY, NO SOLID FOOD!) WATER BLACK COFFEE/TEA (NO CREAMERS, DAIRY/MILK, HONEY) APPLE JUICE, WHITE GRAPE JUICE, LEMONADE POPSCILES, ITALIAN ICE (NOT RED, BLUE, PURPLE) JELL-O (NOT RED, BLUE, PURPLE OR MILK-BASED) GATORADE, PROPEL, POWER-ADE (NOT RED, BLUE, PURPLE)